


# Max/Min Problem

Recall in calculus, given  $f(x)$ , what is maximal/minimal value of  $f(x)$ ?

## Local

To locate critical pts

- $f'(x) = 0$
- or
- $f'(x)$  not defined
- or
- $x$  at boundary



To tell max/min for  $f'(x) = 0$

- $f''(x) > 0$  local min
- $f''(x) < 0$  local max.

Idea: Taylor Expansion of a smooth function

$$f(x) \approx f(x_0) + f'(x_0) \cdot (x - x_0) + \frac{f''(x_0)}{2} (x - x_0)^2$$

For more variable case. given  $f(\vec{x})$   
eg. 2 variables.

$\Rightarrow$

$$f(\vec{x}) \approx f(\vec{x}_0) + \nabla f(\vec{x}_0) \cdot (\vec{x} - \vec{x}_0) +$$

$$\frac{1}{2} \sum_{i,j} \frac{\partial^2 f}{\partial x_i \partial x_j}(\vec{x}_0) (x_i - x_{0,i}) (x_j - x_{0,j})$$

$$\frac{1}{2} [f_{xx}(\vec{x}_0) \cdot x^2 + 2f_{xy}(\vec{x}_0) \cdot xy + f_{yy}(\vec{x}_0) \cdot y^2]$$

## Local

To locate critical pts:

- $\nabla f(\vec{x}) = 0$
- or
- $\nabla f$  is defined
- or
- $\vec{x}$  at boundary.

To tell max/min/saddle

$$A = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{pmatrix}$$

## Global

To locate candidates:

- $f'(x) = 0$
- or
- $f'(x)$  not defined
- or
- $x$  at boundary.

To determine global max/min.

- Compute  $f(x)$  for all candidates

## Global

To locate candidates

- $\nabla f = 0$
- or
- $\nabla f$  not defined
- or
- $\vec{x}$  at boundary.

To determine max/min.

- Compute values for every candidates.

$\det(A) > 0$      $f_{xx} > 0$     local min    

$\det(A) > 0$      $f_{xx} < 0$     local max    

$\det(A) < 0$     saddle pts    

Lagrange Multiplier: Optimization with Constraints.

Goal:  $f(\vec{x})$

(\*) For more constraints

•  $\nabla g = 0$  or  $\nabla h = 0$  or  $\nabla g = k \nabla h$

Constraints:  $g(\vec{x}) = 0$

•  $\nabla f = \lambda_1 \nabla g + \lambda_2 \nabla h$

To locate candidates

•  $\nabla g = 0$

or

•  $\nabla f = \lambda \cdot \nabla g$  for some  $\lambda$ .

} along with  $g(\vec{x}) = 0$

To determine max/min.

- Compute the value for all candidates.

Example: Minimize  $f(x, y) = xy$  subject to the constraint

$$g(x, y) = x^2 + \frac{1}{4}y^2 - 1 = 0$$

Ans. To look for candidates.

$$\textcircled{1} \begin{cases} \nabla g(\vec{x}) = 0 \\ g(\vec{x}) = 0 \end{cases} \Rightarrow \begin{cases} \begin{pmatrix} 2x \\ \frac{1}{2}y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ x^2 + \frac{1}{4}y^2 = 1 \end{cases} \Rightarrow \text{no solution.}$$

$$\textcircled{2} \begin{cases} \lambda \nabla g(\vec{x}) = \nabla f(\vec{x}) \\ g(\vec{x}) = 0 \end{cases} \Rightarrow \begin{cases} \begin{pmatrix} y \\ x \end{pmatrix} = \lambda \cdot \begin{pmatrix} 2x \\ \frac{1}{2}y \end{pmatrix} \\ x^2 + \frac{1}{4}y^2 = 1 \end{cases}$$

$$\begin{cases} y = \lambda \cdot 2x \\ x = \lambda \cdot \frac{1}{2}y \end{cases} \Rightarrow x = \lambda \cdot \frac{1}{2} \cdot \lambda \cdot 2x \Rightarrow x \cdot (\lambda^2 - 1) = 0$$

$$\Rightarrow x = 0 \text{ or } \lambda = \pm 1$$

↓  
impossible.

$$\text{If } \lambda = 1, \quad x^2 + \frac{1}{4} \cdot (2x)^2 = 1 \Rightarrow x = \pm \frac{\sqrt{2}}{2}.$$

$$y = 2x = \pm \sqrt{2}$$

$$\text{If } \lambda = -1, \quad x^2 + \frac{1}{4}(-2x)^2 = 1 \Rightarrow x = \pm \frac{\sqrt{2}}{2}$$

$$y = -2x = \mp \sqrt{2}.$$

So all critical pts are  $(\frac{\sqrt{2}}{2}, \sqrt{2}) \rightarrow f = 1$

$$f = -1 \leftarrow (-\frac{\sqrt{2}}{2}, \sqrt{2}), \quad f = -1 \leftarrow (\frac{\sqrt{2}}{2}, -\sqrt{2}), \quad f = 1 \leftarrow (-\frac{\sqrt{2}}{2}, -\sqrt{2}).$$

Compare the value at all points.

the minimal value is obtained at

$$(\frac{\sqrt{2}}{2}, -\sqrt{2}) \text{ and } (-\frac{\sqrt{2}}{2}, \sqrt{2}), \text{ and is } -1.$$

Example:  $f(x, y) = y^2 - 18x^2 + x^4$

1) What are the local min/max?

$$\nabla f = \begin{pmatrix} -36x + 4x^3 \\ 2y \end{pmatrix} = \vec{0} \Rightarrow \begin{cases} 4x^3 - 36x = 4x \cdot (x^2 - 9) = 0 \\ y = 0 \end{cases} \Rightarrow$$

$x = 0, \pm 3$  and  $y = 0$ . So critical pts are

$$(0, 0) \quad (-3, 0) \quad (3, 0)$$

$$A = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{xy} & f_{yy} \end{pmatrix} = \begin{pmatrix} -36 + 12x^2 & 0 \\ 0 & 2 \end{pmatrix}$$

At  $(0,0)$   $\det(A) = -72 < 0$  saddle pts.


At  $(3,0)$   $\det(A) > 0$   $f_{xx} = 12 \times 9 - 36 = 72 > 0$  local min.

At  $(-3,0)$  local min

2)  $D = [-4, 4] \times [-4, 4]$ . What is the global max/min in  $D$ ?

To locate candidates  $(0,0)$   $(\pm 3,0)$

points on the boundary.



$$f(0,0) = 0 \quad f(\pm 3,0) = 0 - 18 \times 9 + 81 = 81 - 162 = -81$$

$$f(\pm 4, y) = y^2 - 18 \times 4^2 + 4^4 = y^2 + 4^2 \cdot (4^2 - 18) = y^2 - 32$$

$$\text{at most } 16 - 32 = -16 \text{ at } |y| = 4$$

$$\text{at least } 0 - 32 = -32 \text{ at } |y| = 0$$

$$f(x, \pm 4) = 16 - 18x^2 + x^4 = \underbrace{(x^2 - 9)^2 - 81 + 16}$$

$$= (x^2 - 9)^2 - 65$$

$$\text{at most } 9^2 - 65 = 16 \text{ at } |x| = 0$$

$$\text{at least } 7^2 - 65 = -16 \text{ at } |x| = 4$$

Global max is 16 at  $x=0$   $y = \pm 4$

min is -81 at  $x=0$   $y=0$ .